

DATA ASSIMILATION AND MODEL SIMULATIONS IN THE CALIFORNIA CURRENT

Robert L. Haney
Department of Meteorology (MR/Hy)
Naval Postgraduate School
589 Dyer Road
Monterey, CA 93943-5114
phone: (408) 656-4053 fax: (408) 656-3061
e-mail: haneyrl@skimmer.met.nps.navy.mil
Award #: N0001497WR30023

LONG-TERM GOALS

The long-term goal of this research is to develop, verify and apply numerical ocean prediction models to eastern boundary coastal regions in order to improve our scientific understanding of the structure and dynamics of such regions.

OBJECTIVES

The broad objective of this research is to aid in the development of a reliable modeling capability for eastern boundary current regions. The specific objective is to carry out and extensively verify several DieCAST model simulations of the annual cycle in the California Current. We also wish to develop and apply digital filter initialization (DFI) as a diagnostic tool in numerical ocean prediction.

APPROACH

The general approach is to carry out increasingly complex numerical simulations of the California Current using the DieCAST regional model, and to verify the simulations against ONR's extensive eastern boundary current (EBC) data sets. An iterative version of digital filter initialization (DFI) will be verified as a diagnostic tool in several idealized test cases, and then it will be applied to quasi-synoptic hydrographic data from the California Current and the Alboran Sea. Diagnostic results from the Alboran Sea application will be verified using direct measurements of the vertical velocity made by quasi-isobaric RAFOS floats.

WORK COMPLETED

The DieCAST A-grid, low-dispersion (nearly 4th order) regional model (Dietrich 1997) was adapted to the California coastal domain with lateral boundary conditions prescribed using data from both an observed monthly climatology and a global numerical model. Two six-year long numerical simulations were completed. One of the simulations uses a climatological annual cycle of wind stress forcing, while the other simulation includes an

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 1997		2. REPORT TYPE		3. DATES COVERED 00-00-1997 to 00-00-1997	
4. TITLE AND SUBTITLE Data Assimilation and Model Simulations in the California Current				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Postgraduate School, Department of Meteorology, Monterey, CA, 93943				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 4	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

additional wind stress enhancement near each coastal headland that is representative of that observed (eg. Enriquez and Friehe 1995). Preliminary results (below) are very interesting and encouraging.

An iterative form of DFI was applied to the idealized test cases and an optimal version of the method was obtained. It is now being applied to data sets from the California Current (ONR's CTZ and EBC programs) and the Alboran Sea (EC's OMEGA project).

A study of the Atlantic Jet in the Alboran Sea and a numerical simulation of dense water overflow of topography were completed.

RESULTS

Using climatological forcing, the DieCAST model reproduces many of the main features of the observed annual cycle of the California Current including the separation of the coastal jet from the coast in late summer and its offshore migration in autumn and winter. Coastal eddies in the simulation form primarily off the major headlands, especially Cape Mendocino and Point Arena. The surface eddy kinetic energy is in the range of that observed and it undergoes a seasonal cycle with a phase that varies with distance from shore similar to that observed. The additional forcing by the headland wind jets is found to produce both local and remote changes to the simulated annual cycle.

In a separate study, the DieCAST model was shown to simulate dense water overflow of topography better than standard ocean models because of its higher order accuracy and its low numerical dispersion.

The optimal form of DFI is able to diagnose the vertical velocity in an idealized growing baroclinic frontal wave with a relative accuracy of 10%. Using DFI, a closed cyclonic eddy observed off Point Arena California was diagnosed to have negligible vertical velocity on the scale of the eddy itself, but to have very significant vertical velocities ($w \sim 30\text{m/d}$ at 100m) associated with sub-eddy scale meanders in the otherwise circular flow around the eddy. Lagrangian vertical excursions of water parcels was estimated to be over 100m due to the downstream (azimuthal) propagation of the meanders around the eddy. A role for these sub-eddy scale features in the vertical transport of water properties is hypothesized.

Using currents diagnosed by DFI from hydrographic data, the vorticity of the Atlantic Jet in the Alboran Sea was shown to be positive along its entire path, including the part of its path that has anticyclonic (negative) curvature.

IMPACT/APPLICATIONS

Our California Current annual cycle simulations are among the most advanced ones to date. Because of the excellent data sets being used, the model verifications will represent

the standards against which other models will be evaluated in the future. Our results concerning the existence of strong sub-eddy scale vertical velocities off Point Arena are new and, if substantiated by the present OSU research with the EBC data, they suggest a possible role for such features in the vertical eddy transport of water properties.

TRANSITIONS

This research on data assimilation and simulations in the California Current is in broad support of the efforts at FNMOC (M. Clancy) and NRL-Stennis (J. Kindle) to develop a real-time analysis and forecasting capability, including biology, for this region. This support manifests itself in close coordination and timely information exchange on such topics as model properties, experimental set-up, regional modeling difficulties, comparison and exchange of results, methods of model verification, and so forth.

The DFI methodology has been accepted as the major diagnostic tool for estimating the ageostrophic circulation and vertical motion in the EC-sponsored OMEGA project. This multinational European project completed six quasi-synoptic cruises in the western and central Mediterranean last year. DFI will also be utilized by J. Barth (OSU) to study the dynamics of coastal eddies using data from the EBC synoptic surveys.

RELATED PROJECTS

I am collaborating with J. Barth (OSU) in his application of the DFI diagnostic method to the EBC synoptic survey data.

I am collaborating with D. Dietrich (MSU-CAST) in a study of the dense water overflow of topography.

I am collaborating with A. Viudez and J. Tintore (UIB, Spain) in their diagnostic studies of mesoscale variability in the Alboran Sea.

REFERENCES

- Dietrich, D. E., 1997: Application of a modified Arakawa "A" grid ocean model having reduced numerical dispersion to the Gulf of Mexico circulation. *Dyn. Atmos. and Oceans*, (in press).
- Enriquez, A. G., and C. A. Friehe, 1995: Effects of wind stress and wind stress curl variability on coastal upwelling. *J. Phys. Oceanogr.*, **25**, 1651-1671.
- Viudez, A. and R. L. Haney, 1997: On the relative vorticity of the Atlantic Jet in the Alboran Sea. *J. Phys. Oceanogr.*, **27**, 175-185.
- Viudez, A., and R. L. Haney, 1998: The deflection and division of an oceanic baroclinic jet by a coastal boundary. A case study in the Alboran Sea. *J. Phys. Oceanogr.*, **28**, (in press).

- Chumbinho, R., R. L. Haney and S. Ramp, 1997: Kinematics and dynamics of a cyclonic eddy off of Point Arena, California. *J. Phys. Oceanogr.*, (in revision).
- Viudez, A., J. M. Pinot and R. L. Haney, 1997: On the upper layer circulation in the Alboran Sea. *J. Geophys. Res.* (submitted).
- Dietrich, D., R. L. Haney, A. Mehra and S. Piacsek, 1997: Comments on “A Method for Improved Representation of Dense Water Spreading over Topography in Geopotential Coordinate Models”. *J. Phys. Oceanogr.*, (submitted).